

# Numerical Monoids are Totally Cool!<sup>1</sup>

Scott Chapman

*Trinity University, Department of Mathematics, San Antonio, Texas 78212-7200*

---

## Abstract

Several years ago, McDonald's Chicken McNuggets could be purchased only in packs of 6, 9 and 20. Under this restriction, it is not possible to order exactly 43 McNuggets, but there are 2 different ways to order 44 McNuggets:  $4 \cdot 6 + 0 \cdot 9 + 1 \cdot 20 = 44$  and  $1 \cdot 6 + 2 \cdot 9 + 1 \cdot 20 = 44$ . This computation is a modern version of an older problem commonly known as the *postage stamp problem*.

The ideas in the first paragraph can be mathematically formalized as follows. Let  $n_1, n_2, \dots, n_t$  be a finite set of positive integers. Then the numerical monoid  $S$  generated by  $\langle n_1, n_2, \dots, n_t \rangle$  is the set of all  $s$  such that  $s = \sum a_i n_i$ , where each  $a_i$  is a nonnegative integer. If the integers  $n_1, n_2, \dots, n_t$  are relatively prime, then it can be shown that there is a largest integer  $m$  such that  $m \notin S$ . This  $m$  is known as the *Frobenius number* of  $S$  and because of the difficulty in its computation, has a long and elusive mathematical history. In addition to discussing the Frobenius number, we shall also talk about how elements factor in a numerical monoid. For  $s = \sum a_i n_i$  in  $S$ ,  $\sum a_i$  is called a *factorization length* of  $s$ . Let  $\mathcal{L}(s) = \{m_1, m_2, \dots, m_k\}$  (where  $m_i < m_{i+1}$  for  $1 \leq i \leq k-1$ ) be the set of all possible factorization lengths of  $s$ . The sets  $\mathcal{L}(s)$  have interesting properties, and I will discuss in some details results concerning these sets obtained by my past REU students.

---

---

*Email address:* [schapman@trinity.edu](mailto:schapman@trinity.edu) (Scott Chapman).

<sup>1</sup> The title for this talk is a quote from one of my former REU students Nathan Kaplan who is currently a Senior at Princeton.