

Extremals for the families of commuting  
spherical contractions and their adjoints.

Stefan Richter

joint work with Carl Sundberg

University of Tennessee

April 2009

Topic: Dilations and extensions of  $d$ -tuples of commuting Hilbert space operators

Example for  $d = 1$ :

**Thm 1.** (*the Sz.-Nagy dilation theorem*)

$$T \in \mathcal{B}(\mathcal{H}), \|T\| \leq 1$$

$$\Rightarrow \exists V \in \mathcal{B}(\mathcal{K}), \mathcal{H} \subseteq \mathcal{K},$$

$$V\mathcal{H} \subseteq \mathcal{H}, \|V^*x\| = \|x\|,$$

$$T = V|_{\mathcal{H}}.$$

$V =$  co-isometric extension of  $T$

$$V = S^* \oplus U,$$

$S$  unilateral shift of some multiplicity,

$U$  unitary

A study of such  $V$  leads to function theory in  $\mathbb{D}$ .

All Hilbert spaces in the following are supposed to be separable.

$$d \in \mathbb{N}, \quad \mathbb{B}^d = \{z \in \mathbb{C}^d : |z| < 1\}$$

**Defn 2. (Agler)** A family  $\mathcal{F}$  is a collection of  $d$ -tuples  $T = (T_1, \dots, T_d)$  of Hilbert space operators,  $T_i \in \mathcal{B}(\mathcal{H})$  such that

(a)  $\mathcal{F}$  is bounded,

$$\exists c > 0 \quad \forall T = (T_1, \dots, T_d) \in \mathcal{F} : \|T_i\| \leq c \quad \forall i$$

(b) restrictions to invariant subspaces

$$T \in \mathcal{F}, \mathcal{M} \subseteq \mathcal{H}, T_i \mathcal{M} \subseteq \mathcal{M} \quad \forall i \Rightarrow T|_{\mathcal{M}} \in \mathcal{F}$$

(c) direct sums

$$T_n \in \mathcal{F} \Rightarrow \bigoplus_n T_n \in \mathcal{F}, \quad T_n = (T_{1n}, \dots, T_{dn})$$

(d) unital  $*$ -representations

$$\pi : \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{K}), \pi(I) = I,$$

$$T = (T_1, \dots, T_d) \in \mathcal{F}$$

$$\Rightarrow \pi(T) = (\pi(T_1), \dots, \pi(T_d)) \in \mathcal{F}.$$

Examples:

$d = 1$  :

$\mathcal{F} =$  contractions,  $T^*T \leq I$

isometries,  $T^*T = I$

subnormal contractions

$d \geq 1$  :

$\mathcal{F} =$  contractions

commuting contractions

isometries

commuting isometries

$\mathcal{F} =$  commuting spherical contractions

commuting row contractions (d-contractions)

commuting spherical isometries

**Defn 3.** If  $T = (T_1, \dots, T_d), T_i \in \mathcal{B}(\mathcal{H}),$   
 $S = (S_1, \dots, S_d), S_i \in \mathcal{B}(\mathcal{K}), T, S \in \mathcal{F},$   
then

$$\begin{aligned} T \leq S &\Leftrightarrow \mathcal{H} \subseteq \mathcal{K}, S\mathcal{H} \subseteq \mathcal{H}, T = S|_{\mathcal{H}} \\ &\Leftrightarrow S = \begin{pmatrix} T & X \\ 0 & Y \end{pmatrix} \\ &\Leftrightarrow S \text{ extends } T. \end{aligned}$$

**Defn 4.**  $T$  is **extremal** for  $\mathcal{F},$

$$\Leftrightarrow S \geq T, S \in \mathcal{F} \Rightarrow S = \begin{pmatrix} T & 0 \\ 0 & Y \end{pmatrix} = T \oplus Y.$$

We will write  $T \in \text{ext}(\mathcal{F})$

**Thm 5.** (Agler)

$\mathcal{F}$  family,  $T \in \mathcal{F} \Rightarrow \exists S \in \text{ext}(\mathcal{F}) S \geq T$

## Examples:

$\mathcal{F} = \text{contractions} \Rightarrow \text{ext}(\mathcal{F}) = \text{co-isometries}$

**Cor 6.** (*Sz. Nagy*)

*Every contraction has a co-isometric extension.*

$\mathcal{F} = \text{isometries} \Rightarrow \text{ext}(\mathcal{F}) = \text{unitaries}$

**Cor 7.** *Every isometry is subnormal.*

$\mathcal{F} = \text{subnormal contractions}$

$\Rightarrow \text{ext}(\mathcal{F}) = \text{normal contractions.}$

**Cor 8.** *Every subnormal contraction is subnormal.*

Commuting spherical isometries (A. Athavale)

$$\mathcal{F} = \{T = (T_1, \dots, T_d) : T_i \leftrightarrow T_j, \\ \sum_{i=1}^d \|T_i x\|^2 = \|x\|^2 \quad \forall x \in \mathcal{H}\}$$

$$\text{ext}(\mathcal{F}) = \{U = (U_1, \dots, U_d) : U_i \leftrightarrow U_j, U_i \text{ normal} \\ \sum_{i=1}^d \|U_i x\|^2 = \|x\|^2 \quad \forall x \in \mathcal{H}\}$$

= commuting spherical unitaries

The proof follows Attele-Lubin, JFA, 1996.

**Cor 9.** (Athavale, 91) *Every commuting spherical isometry is jointly subnormal.*

## Commuting spherical contractions

Drury 78, Mueller-Vasilescu 93,  
Arveson 98

$$\mathcal{F} = \{T = (T_1, \dots, T_d) : T_i \leftrightarrow T_j, \\ \sum_{i=1}^d \|T_i x\|^2 \leq \|x\|^2 \quad \forall x \in \mathcal{H}\}$$

$$T \in \mathcal{F} \Leftrightarrow \sum_{i=1}^d T_i^* T_i \leq I, \text{ commuting}$$

$$\text{ext}(\mathcal{F}) = \{S^* \oplus U\}$$

$U =$  commuting spherical unitary

$S = d$ -shift of some multiplicity

$$S = M_z \text{ on } H_d^2(\mathcal{D}) = H_d^2 \otimes \mathcal{D}$$

$H_d^2 \subseteq \text{Hol}(\mathbb{B}^d)$ , Drury-Arveson-Hardy space  
defined by reproducing kernel

$$k_w(z) = \frac{1}{1 - \langle z, w \rangle}, \quad z, w \in \mathbb{B}^d$$

**Thm 10.** (Richter-Sundberg) Let  $T = (T_1, \dots, T_d)$  be a commuting operator tuple.

Then the following are equivalent

(a)  $T \in \text{ext}(\mathcal{F})$

(b)  $T = S^* \oplus U$

(c) (1)  $\sum_{i=1}^d T_i^* T_i = P = \text{a projection}$

(2)  $\sum_{i=1}^d T_i T_i^* \geq I$

(3) If  $x_1, \dots, x_d \in \mathcal{H}$  with  $T_i x_j = T_j x_i$ , then  $\exists x \in \mathcal{H}$  with  $x_i = T_i x$ .

(c3) says that the Koszul complex for  $T$  is exact at a certain stage.

Note 1: For  $d = 1$  (c) becomes

(1)  $T^*T = P$ , i.e.  $T$  is a partial isometry

(2)  $TT^* \geq I$ , so  $T$  is onto

(3) if  $x_1 \in \mathcal{H}$ , then  $\exists x \in \mathcal{H}$  with  $x_1 = Tx$ ,  
i.e.  $T$  is onto

Hence (1)&(2) or (1)&(3) are equivalent to  $T^*$  being an isometry.

For  $d > 1$  let  $T = M_z$  on  $H^2(\partial\mathbb{B}^d)$ , then (1) and (3) are satisfied, but (2) is not.

Note 2: If  $T \in \mathcal{F}$  and

if  $\sum_{i=1}^d T_i^* T_i \neq$  a projection, then  $T \notin \text{ext}(\mathcal{F})$ .

Commuting row contractions (d-contractions)

$$\mathcal{F} = \{T : T^* \text{ is a commuting spherical contraction} \}$$

$$= \{T : T_i \leftrightarrow T_j, \sum_{i=1}^d \|T_i^* x\|^2 \leq \|x\|^2 \forall x\}$$

$$= \{T : T_i \leftrightarrow T_j, \|\sum_{i=1}^d T_i x_i\|^2 \leq \sum_{i=1}^d \|x_i\|^2 \quad \forall x_i\}$$

$$T \in \mathcal{F}$$

$$\Leftrightarrow (T_1, \dots, T_d) : \mathcal{H} \oplus \dots \oplus \mathcal{H} \rightarrow \mathcal{H} \text{ is contractive} \\ \text{commutative}$$

$$\Rightarrow T^* = S^* \oplus U|_{\mathcal{H}}$$

(by Mueller/Vasilescu-Arveson)

$$\Rightarrow T = P_{\mathcal{H}}(S \oplus U^*)|_{\mathcal{H}},$$

$\mathcal{H}$  =co-invariant for  $S \oplus U^*$

$$\text{ext}(\mathcal{F}) = ?$$

$$D_* = \left( I - \sum_{i=1}^d T_i T_i^* \right)^{1/2}$$

**Thm 11.** (easy)

(a) If  $D_* = 0$ , then  $T \in \text{ext}(\mathcal{F})$

spherical co-isometries

(b) If  $D_*$  is onto, then  $T \notin \text{ext}(\mathcal{F})$ .

Thus, if  $\exists c < 1 \ni \sum_i \|T_i^* x\|^2 \leq c \|x\|^2$ , then  $T \notin \text{ext}(\mathcal{F})$ .

(c) If  $D_*$  is a projection, then

$T \notin \text{ext}(\mathcal{F})$

$$\Leftrightarrow \exists x_1, \dots, x_d \in \text{ran } D_*, \sum_{i=1}^d \|x_i\|^2 > 0$$

with  $T_i x_j = T_j x_i$

If  $S = M_z = d$ -shift, then

$$D_*^2 = I - \sum_{i=1}^d S_i S_i^*$$

is a rank 1 projection, and

$$\text{ran } D_* = \text{constants},$$

hence  $S \in \text{ext}(\mathcal{F})$ .

**Cor 12.** If  $d > 1$ , then  $\{S \oplus U^*\} \subsetneq \text{ext}(\mathcal{F})$

$U = \text{spherical unitary}$

Recall

$$\{S^* \oplus U\} = \text{ext}(\mathcal{F}^*)$$

Example for  $\neq$ :

$$T = (M_z^*, H^2(\partial\mathbb{B}^d)) \in \text{ext}(\mathcal{F}).$$

$$D_* = (I - \sum_{i=1}^d T_i T_i^*)^{1/2}$$

**Thm 13. (R-S)**

*If  $T \in \mathcal{F}$  and if  $D_*$  has rank one, i.e.*

$$D_* = u \otimes u$$

*for some  $u \neq 0$ , then*

$$T \in \text{ext}(\mathcal{F}) \Leftrightarrow \dim \text{span}\{u, T_1 u, \dots, T_d u\} \geq 3$$

If  $S = (M_z, H_d^2) =$  the  $d$ -shift,  
if  $\mathcal{M}$  is invariant for  $S$ ,  $\mathcal{M} \neq H_d^2$   
then

$$T = P_{\mathcal{M}^\perp} S|_{\mathcal{M}^\perp} \in \mathcal{F},$$

and  $D_*$  has rank 1:

$$\begin{aligned} D_*^2 &= I_{\mathcal{M}^\perp} - \sum_{i=1}^d P_{\mathcal{M}^\perp} S_i S_i^* P_{\mathcal{M}^\perp} \\ &= P_{\mathcal{M}^\perp} (I - \sum S_i S_i^*) P_{\mathcal{M}^\perp} \\ &= u \otimes u, \end{aligned}$$

$$u = P_{\mathcal{M}^\perp} \mathbf{1} \neq 0$$

**Cor 14.** *If  $\mathcal{M} \neq H_d^2$ , if*

$$\mathcal{L} = \left\{ a + \sum_{i=1}^d b_i z_i \right\},$$

*then  $T = P_{\mathcal{M}^\perp} S|_{\mathcal{M}^\perp} \notin \text{ext}(\mathcal{F})$  if and only if*

$$\dim \mathcal{M} \cap \mathcal{L} \in \{d-1, d\}$$

In fact, in this case, if  $T \notin \text{ext}(\mathcal{F})$ , then

$$T_i = a_i I + b_i S$$

for some  $S$  with  $\text{rank}(I - SS^*) = 1$ .

The Theorem can be used to produce examples

$$T \in \text{ext}(\mathcal{F}) \text{ but } D_* \neq \text{a projection.}$$

Thus THM 11(c) does not not characterize all extremals.