

DIFFERENTIAL HARNACK INEQUALITIES VIA CONCAVITY OF THE ARRIVAL TIME

THEODORA BOURNI AND MAT LANGFORD

ABSTRACT. We present a simple connection between differential Harnack inequalities for hypersurface flows and natural concavity properties of their time-of-arrival functions. We prove these concavity properties directly for a large class of flows by applying a concavity maximum principle argument to the corresponding level set flow equations. In particular, this yields a short proof of Hamilton's differential Harnack inequality for mean curvature flow and, more generally, Andrews' differential Harnack inequalities for certain " α -inverse-concave" flows.

1. CONCAVITY MAXIMUM PRINCIPLES

Our goal is to deduce concavity properties for the time-of-arrival functions of a large class of geometric flow equations using the concavity maximum principle. The main idea, due to Korevaar [20] and later extended by Kennington [19] and Kawohl [18] is summarized in the following theorem.

Theorem 1.1. *Let $\Omega \subset \mathbb{R}^n$ be a bounded, convex, open set and suppose that $u \in C^1(\overline{\Omega})$ is twice differentiable in Ω and satisfies the equation*

$$-f(Du(x), D^2u(x)) = b(x, u(x), Du(x)) \text{ in } \Omega$$

with $f : \mathbb{R}^n \times \Gamma \rightarrow \mathbb{R}$, $\Gamma \underset{\text{convex, open}}{\subset} \text{Sym}^{n \times n}$, satisfying

(i) *Weak ellipticity:*

$$r \geq s \implies f(p, r) \geq f(p, s).$$

(ii) *Concavity:*

$$f(p, \lambda r + (1 - \lambda)s) \geq \lambda f(p, r) + (1 - \lambda)f(p, s).$$

and $b : \Omega \times \mathbb{R} \times \mathbb{R}^n \rightarrow \mathbb{R}$ satisfying

(iii) *Monotonicity:*

$$z > w \implies b(x, z, p) < b(x, w, p).$$

(iv) *Joint concavity:*

$$b(\lambda(x, z) + (1 - \lambda)(y, w), p) \geq \lambda b(x, z, p) + (1 - \lambda)b(y, w, p).$$

If the graph of u lies below its boundary tangent hyperplanes, then u is concave.

Proof. The argument is essentially that of Korevaar [20]: Consider Korevaar's "concavity function" $Z : [0, 1] \times \Omega \times \Omega \rightarrow \mathbb{R}$, defined by [20]

$$(1) \quad Z(r, x, y) := u(rx + (1-r)y) - (ru(x) + (1-r)u(y)).$$

This function measures how far the point $(rx + (1-r)y, u(rx + (1-r)y))$ in $\bar{\Omega} \times \mathbb{R}$ lies above the line joining the points $(x, u(x))$ and $(y, u(y))$. We need to prove that $Z \geq 0$.

Choose the triple (r_0, x_0, y_0) so that

$$Z(r_0, x_0, y_0) = \min_{[0,1] \times \bar{\Omega} \times \bar{\Omega}} Z(r, x, y).$$

If $r_0 = 0$ or $r_0 = 1$, then $Z(r_0, x_0, y_0) = 0$, which implies the claim. So we may assume that $r_0 \in (0, 1)$. Suppose that $x_0 \in \partial\Omega$. If $Z(r_0, x_0, y_0) < 0$, then, since the graph of u lies below its boundary tangent hyperplanes, it would be possible to find a point (r_1, x_1, y_1) with $Z(r_1, x_1, y_1) < Z(r_0, x_0, y_0)$ by moving x_0 a small amount inwards along the line joining x_0 and y_0 , contradicting minimality of (r_0, x_0, y_0) [20]. Indeed, consider the function

$$f(\varepsilon) := Z(x_\varepsilon, y_0, r_\varepsilon),$$

where $x_\varepsilon := x_0 + \varepsilon(y_0 - x_0)$ and $r_\varepsilon := r_0/(1 - \varepsilon)$. Since

$$r_\varepsilon x_\varepsilon + (1 - r_\varepsilon)y_0 \equiv z_0,$$

the boundary condition implies that

$$\left. \frac{d}{d\varepsilon} f \right|_{\varepsilon=0} = -r_0(u(y_0) - u(x_0) - Du(x_0) \cdot (y_0 - x_0)) < 0,$$

in contradiction with the fact that $f(\varepsilon)$ is minimized at $\varepsilon = 0$.

A similar argument applies at y_0 . So we may assume that x_0 and y_0 are interior points.

Let us abuse notation by writing $Z(x, y) := Z(r_0, x, y)$. Then (x_0, y_0) is a stationary point of Z and hence, setting $z_0 := r_0 x_0 + (1 - r_0)y_0$,

$$(2a) \quad 0 = \partial_{x^i} Z(x_0, y_0) = r_0(u_i(z_0) - u_i(x_0))$$

and

$$(2b) \quad 0 = \partial_{y^i} Z(x_0, y_0) = (1 - r_0)(u_i(z_0) - u_i(y_0)).$$

So

$$(3) \quad Du(z_0) = Du(x_0) = Du(y_0) =: p_0.$$

Since (x_0, y_0) is a local minimum,

$$\begin{aligned} 0 &\leq (\partial_{x^i} + \partial_{y^i})(\partial_{x^j} + \partial_{y^j})Z(x_0, y_0) \\ &= u_{ij}(z_0) - r_0 u_{ij}(x_0) - (1 - r_0)u_{ij}(y_0). \end{aligned}$$

The ellipticity and concavity of f and the joint-concavity of b then imply

$$\begin{aligned} b(z_0, u(z_0), p_0) &= -f(p_0, D^2u(z_0)) \\ &\leq -f(p_0, r_0 D^2u(x_0) + (1 - r_0)D^2u(y_0)) \\ &\leq -r_0 f(p_0, D^2u(x_0)) - (1 - r_0)f(p_0, D^2u(y_0)) \\ &= r_0 b(x_0, u(x_0), p_0) + (1 - r_0)b(y_0, u(y_0), p_0) \\ &\leq b(z_0, r_0 u(x_0) + (1 - r_0)u(y_0), p_0). \end{aligned}$$

The claim now follows from the monotonicity of b . \square

Remark 1.2. *Note that in Theorem 1.1, although the solution u is required to be twice differential in Ω and C^1 up to the boundary, no regularity hypotheses are needed for the functions f and b .*

Remark 1.3. *In the quasi-linear setting, Theorem 1.1 recovers the original result of Korevaar [20]. It also immediately recovers a special case of an important refinement observed by Kennington [19] (see also Kawohl [18, Theorem 3.13]). Indeed, if $b > 0$, then we may rewrite the quasi-linear equation*

$$-a^{ij}(Du(x))u_{ij}(x) = b(x, u(x), Du(x))$$

as

$$-f(Du(x), D^2u(x)) = b_*(x, u(x), Du(x)),$$

where

$$f(p, r) := -(a^{ij}(p)r_{ij})^{-1} \quad \text{and} \quad b_*(x, z, p) := -b^{-1}(x, z, p).$$

If a and b satisfy the conditions of [19, Theorem 3.1] (see also [18, Theorem 3.13]), then f is weakly elliptic and concave, and b_ is decreasing and joint-concave. So the equation is of the form allowed by Theorem 1.1. It is worth noting that, by allowing the left hand side to depend nonlinearly on the second derivatives, the proof is actually simplified compared to the arguments presented in [18] and [19].*

Remark 1.4. *Theorem 1.1 can be almost immediately applied (cf. [20]) to the nonlinear capillary problem*

$$(4) \quad \begin{aligned} F(A[u]) &= \kappa u + \lambda \quad \text{in } \Omega \\ \nu|_{\text{graph } u} &= \nu|_{\Omega} \quad \text{on } \partial\Omega, \end{aligned}$$

where $\kappa > 0$ and F is a non-decreasing, concave function of the second fundamental form $A[u]$ of graph u . Indeed, the function $v := C - u$, $C > \max_{\overline{\Omega}} u$, satisfies an equation of the form

$$\begin{aligned} -f(Dv, D^2v) &= \kappa v^{-1} \text{ in } \Omega \\ \nu|_{\text{graph } v} &= \nu|_{\Omega} \text{ on } \partial\Omega, \end{aligned}$$

where f is nondecreasing and concave in its second argument. Although Dv blows-up at the boundary of Ω , Theorem 1.1 applies when we restrict to domains $\Omega' \Subset \Omega$ which are sufficiently close to Ω . So the conclusion holds in all $\Omega' \Subset \Omega$ and we conclude that u is convex in Ω .

In some cases, a perturbation argument (cf. [20, Lemma 1.5]) can be used to weaken Condition (iii) to weak monotonicity, in which case the theorem can be used to study certain nonlinear Weingarten problems ($\kappa = 0$ in (4)) and certain nonlinear eigenvalue problems. We will not explore such applications here, since they have been developed elsewhere (see [1, 17, 18, 19, 20, 22]).

In Section 3, we apply this simple and elegant idea to certain degenerate fully nonlinear equations (namely, level set flows of convex hypersurfaces).

Let us begin our investigation within the simpler context of *mean curvature flow*, where our main result follows more or less as in Theorem 1.1. (A more subtle argument will be required when we consider more general flows.)

2. MEAN CURVATURE FLOW

Let $\{\mathcal{M}_t^n\}_{t \in [t_0, T]}$ be a family of smooth, strictly convex boundaries $\mathcal{M}_t^n = \partial\Omega_t$ moving with normal velocity $-H\nu$, where $\nu(x, t)$ is the outward pointing unit normal to \mathcal{M}_t^n at x and $H = \text{div } \nu$ is the corresponding mean curvature. Recall that the *arrival time* $u : \cup_{t \in [t_0, T]} \mathcal{M}_t^n \rightarrow \mathbb{R}$ of the family $\{\mathcal{M}_t^n\}_{t \in [t_0, T]}$ is defined by

$$u(p) = t \iff p \in \mathcal{M}_t^n.$$

Note that u is well-defined since the hypersurfaces move monotonically.

Let $X : M^n \times [t_0, T] \rightarrow \mathbb{R}^{n+1}$ be a smooth family of parametrizations $X(\cdot, t)$ of \mathcal{M}_t^n . Then

$$(5) \quad u(X(x, t)) = t.$$

Fix a point $q = X(x, t)$ in \mathcal{M}_t^n and local orthonormal coordinates $\{x^i\}_{i=1}^n$ for M^n about x (with respect to the induced metric at time

t). Choose the basis $\{e_i\}_{i=1}^{n+1}$ for \mathbb{R}^{n+1} so that $e_{n+1} = \nu(x, t)$ and $e_i = \partial_i X(x, t)$ for each $i = 1, \dots, n$. Differentiating (5) yields the identities

$$(6) \quad Du \cdot \partial_i X = 0 \quad \text{and} \quad -H Du \cdot \nu = 1$$

and hence

$$(7) \quad Du = -\frac{\nu}{H}.$$

Since $H = \operatorname{div} \nu$, we deduce that u satisfies the *level set (mean curvature) flow*

$$(8) \quad -|Du| \operatorname{div} \left(\frac{Du}{|Du|} \right) = 1.$$

Moreover, differentiating (6) at the point (x, t) , we obtain

$$(9) \quad D^2 u = \begin{pmatrix} -A/H & \nabla H/H^2 \\ \nabla H/H^2 & -\partial_t H/H^3 \end{pmatrix}.$$

It follows that $w := \sqrt{2(u - t_0)}$ satisfies

$$(10) \quad D^2 w = w^{-1} \begin{pmatrix} -A/H & \nabla H/H^2 \\ \nabla H/H^2 & -(\partial_t H + H/w^2)/H^3 \end{pmatrix}.$$

This is equivalent to the bilinear form studied by Hamilton in his derivation of the differential Harnack inequality [13] (and later by Chow–Chu [10], Kotschwar [21], and Helmsendorfer–Topping [14], who formulated “space-time” approaches to differential Harnack inequalities).

Recall that the differential Harnack inequality asserts that

$$(11) \quad \partial_t H + 2\nabla_V H + A(V, V) + \frac{H}{2(t - t_0)} \geq 0 \quad \text{for all } V \in T\mathcal{M}_t, \quad t > t_0.$$

It is easy to see that local concavity of w is equivalent to (11): Fix $p \in \mathcal{M}_t^n$ and any $V \in T_p \mathbb{R}^{n+1}$. Then, either V is tangent to \mathcal{M}_t^n , in which case

$$w D^2 w(V, V) = -\frac{A(V, V)}{H},$$

or $V = \lambda(V^\top - H\nu)$ for some $\lambda \in \mathbb{R}$ and $V^\top \in T_p \mathcal{M}_t^n$, in which case

$$(12) \quad w D^2 w(V, V) = -\frac{\lambda^2}{H} \left(\partial_t H + \frac{H}{2(t - t_0)} + 2\nabla_{V^\top} H + A(V^\top, V^\top) \right).$$

Since the Harnack inequality is saturated by self-similarly expanding solutions, so is local concavity of the square root of the arrival time. In fact, this is readily deduced directly: if $\mathcal{M}_t^n = \sqrt{t} \mathcal{M}_1^n$, for $t > 0$, defines a self-similarly expanding solution, then $w = u^{\frac{1}{2}}$ is homogeneous of degree 1 since $\sqrt{t/s} X \in \mathcal{M}_t^n$ if and only if $X \in \mathcal{M}_s^n$. But then $D^2 w$ is degenerate in radial directions.

For *ancient* solutions $\{\mathcal{M}_t^n\}_{t \in (-\infty, T)}$, the Harnack inequality becomes

$$(13) \quad \partial_t H + 2\nabla_V H + A(V, V) \geq 0 \text{ for all } V \in T\mathcal{M}_t, t > -\infty,$$

which, by the same argument, is seen to be equivalent to local concavity of u itself.

Theorem 2.1. *Let $\Omega \subset \mathbb{R}^{n+1}$ be a bounded, convex, open set with smooth boundary. Given $u_0 \in \mathbb{R}$, suppose that $u \in C^1(\overline{\Omega})$ has a single critical point, $p \in \Omega$, is twice differentiable in $\Omega \setminus \{p\}$, and satisfies*

$$(14) \quad \begin{cases} -|Du| \operatorname{div} \left(\frac{Du}{|Du|} \right) = 1 & \text{in } \Omega \setminus \{p\} \\ u \equiv u_0 & \text{on } \partial\Omega. \end{cases}$$

Then $\sqrt{2(u - u_0)}$ is concave.

If we no longer assume that Ω is bounded, but require instead that $u_0 = -\infty$ and that the level sets of u are bounded and convex, then u is concave.

Proof. Set $w := \sqrt{2(u - u_0)}$. Then

$$Dw = \frac{Du}{w}$$

and hence

$$-\sum_{i,j=1}^n \left(\delta_{ij} - \frac{w_i w_j}{|Dw|^2} \right) w_{ij} = -|Dw| \operatorname{div} \left(\frac{Dw}{|Dw|} \right) = w^{-1}$$

in $\Omega \setminus \{p\}$. Observe that the tangent hyperplanes to the graph of w are vertical at the boundary. Indeed, the normal to the graph of w is given by

$$N = \frac{(-Dw, 1)}{\sqrt{1 + |Dw|^2}} = \frac{(\nu, Hw)}{\sqrt{H^2 w^2 + 1}}.$$

The concavity maximum principle now implies that w is concave. We proceed as in the proof of Theorem 1.1: Let (r_0, x_0, y_0) attain the minimum of the concavity function Z (defined in (1)). Since graph w lies below its boundary tangent hyperplanes (see Remark 1.4), we may assume that (r_0, x_0, y_0) is an interior point. So we obtain the gradient identities (2a)-(2b) and hence $p_0 := Du(x_0) = Du(y_0)$. If p_0 is not zero, the argument given in Theorem 1.1 implies that $Z(r_0, x_0, y_0) \geq 0$. On the other hand, if $p_0 = 0$, then $x_0 = y_0$ (since, by hypothesis, u has but one critical point) and hence $Z(r_0, x_0, y_0) = 0$.

To prove the second claim, fix any point $p \in \Omega$ and any $t < u(p)$. Then $p \in \Omega_t := \{q \in \Omega : u(q) > t\}$. The hypotheses on u imply that

Ω_t is bounded and hence, by the first part of the theorem, the function $w : \Omega_t \rightarrow \mathbb{R}$ given by $w(q) = (2(u(q) - t))^{\frac{1}{2}}$ is concave. Thus,

$$D^2u(p) = w^{-1}(p)D^2w(p) + \frac{Du(p) \otimes Du(p)}{w^2(p)} \leq \frac{Du(p) \otimes Du(p)}{2(u(p) - t)}.$$

Taking $t \rightarrow -\infty$ yields the claim. \square

Note that, for an initial hypersurface which bounds a bounded convex body, the corresponding solution to mean curvature flow remains smooth until it contracts to a single point, p say. It follows that the arrival time is smooth away from its only critical point, p , and C^1 at p . In fact, Huisken [15] proved that the solution becomes ‘asymptotically round’ near p , which actually implies that the arrival time is of class¹ C^2 [16]. In any case, Theorem 2.1 provides a rather simple proof of Hamilton’s differential Harnack inequality.

Corollary 2.2. *Let $\{\mathcal{M}_t^n\}_{t \in [t_0, T]}$ be a smooth family of boundaries $\mathcal{M}_t^n = \partial\Omega_t$ of bounded convex bodies Ω_t evolving by mean curvature. Suppose that the boundaries \mathcal{M}_t^n contract to a point at time T . Then the square root $w := (2(u - t_0))^{\frac{1}{2}}$ of the arrival time $u : \Omega_{t_0} \rightarrow \mathbb{R}$ is concave. Equivalently,*

$$\partial_t H + 2\nabla_V H + A(V, V) + \frac{H}{2(t - t_0)} \geq 0 \text{ for all } V \in T\mathcal{M}_t, \quad t \in (t_0, T).$$

If the solution is ancient, then u is concave. Equivalently,

$$\partial_t H + 2\nabla_V H + A(V, V) \geq 0 \text{ for all } V \in T\mathcal{M}_t, \quad t \in (t_0, T).$$

Proof. By (7), we find that u has a single critical point and is differentiable everywhere. It follows that the arrival time u of the family satisfies the hypotheses of Theorem 2.1, and we conclude that its square root $w := (2(u - t_0))^{\frac{1}{2}}$ is concave. The differential Harnack inequality then follows from (10) as in (12). The remaining claim is proved similarly. \square

Remark 2.3. *Note that, since the level-set flow equation is not defined when $Du = 0$, a separate argument in Theorem 2.1 was necessary at such points.*

After we completed this work, we learned that Trudinger had essentially pointed out the proof of Theorem 2.1 in the concluding remarks to [23], and that Evans and Spruck [12, Theorem 7.6] had proved a

¹Colding and Minicozzi [11] proved that the arrival time of a general compact, mean convex mean curvature flow is twice differentiable. But this result requires the full force of the structure theory for singularities in mean curvature flow. We only require here that the hypersurfaces shrink to a (not necessarily round) point.

stronger version of Theorem 2.1 by applying the concavity maximum principle to approximating solutions to the (non-singular) ε -regularized level-set flow and taking a limit as $\varepsilon \rightarrow 0$. Notably, both of these works preceded Hamilton's paper [13].

Xu-Jia Wang [24, Lemma 4.1] observed that the logarithm of $u - t_0$ is concave (in general), and used this to deduce that u is concave for an ancient solution. His argument seems to implicitly make use of the assumptions in Theorem 2.1 and was one of the motivations for this work.

3. FLOWS BY NONLINEAR FUNCTIONS OF CURVATURE

We now consider a much larger class of evolutions. Let $\{\mathcal{M}_t^n\}_{t \in [t_0, T]}$ be a family of smooth, convex boundaries $\mathcal{M}_t^n = \partial\Omega_t$ moving with normal velocity $-F\nu$, where $\nu(x, t)$ is the outward pointing unit normal to \mathcal{M}_t^n at x . We consider speeds $F(\cdot, t) : \mathcal{M}_t^n \rightarrow \mathbb{R}$ given by

$$(15) \quad F(x, t) = f^\alpha(\nu(x, t), [A_{(x,t)}])$$

for some $\alpha > 0$, where $A_{(x,t)}$ is the second fundamental form of \mathcal{M}_t^n at x and $[A_{(x,t)}]$ its component matrix with respect to an orthonormal frame for $T_x\mathcal{M}_t^n$, and $f : S^n \times \Gamma_+^{n \times n} \rightarrow \mathbb{R}$ is a smooth, positive function which is $\text{SO}(n)$ -invariant² and monotone non-decreasing in its second entry, where $\Gamma_+^{n \times n}$ is the cone of positive definite, symmetric $n \times n$ matrices.

Since f is positive, the hypersurfaces move monotonically inwards, so the arrival time $u : \cup_{t \in [t_0, T]} \mathcal{M}_t^n \rightarrow \mathbb{R}$, which we recall is given by

$$u(p) = t \iff p \in \mathcal{M}_t^n,$$

is well-defined. If the boundaries contract to a point, then the arrival time is well-defined on all of Ω_{t_0} and of class $C^1(\overline{\Omega})$. If F is isotropic and the boundaries contract smoothly to a 'round' point, then the arrival time is of class $C^2(\overline{\Omega})$. Indeed, the same calculations as in the preceding section reveal that

$$(16) \quad Du = -\frac{\nu}{F}$$

and

$$(17) \quad D^2u = \begin{pmatrix} -A/F & \nabla F/F^2 \\ \nabla F/F^2 & -\partial_t F/F^3 \end{pmatrix}.$$

Since, in the isotropic case,

$$\partial_t F = \dot{F}(\nabla^2 F + FA^2),$$

²I.e. invariant under conjugation of its second factor by special orthogonal matrices.

where $\dot{F} := Df|_{[A]}$, the claims follow similarly as in [16]. Moreover, u satisfies the *level set flow*

$$|Du|f^\alpha \left(-\frac{Du}{|Du|}, -D\frac{Du}{|Du|} \right) = 1.$$

Set

$$w := ((1 + \alpha)(u - t_0))^{\frac{1}{1+\alpha}}.$$

Then, away from the final point,

$$Dw = w^{-\alpha} Du,$$

$$\begin{aligned} (18) \quad w^\alpha D^2 w &= D^2 u - \alpha \frac{Du \otimes Du}{w^{1+\alpha}} \\ &= \begin{pmatrix} -A/F & \nabla F/F^2 \\ \nabla F/F^2 & -(\partial_t F + \alpha F/w^{1+\alpha})/F^3 \end{pmatrix} \end{aligned}$$

and

$$|Dw|f^\alpha \left(-\frac{Dw}{|Dw|}, -\frac{1}{|Dw|} \left[I - \frac{Dw \otimes Dw}{|Dw|^2} \right] \cdot D^2 w \right) = w^{-\alpha}.$$

As in [5], let us call a function $f : S^n \times \Gamma_+^{n \times n} \rightarrow \mathbb{R}$ *inverse-concave* if the dual function $f_* : S^n \times \Gamma_+^{n \times n} \rightarrow \mathbb{R}$ defined by

$$f_*^{-1}(p, r) = f(p, r^{-1})$$

is concave.

Theorem 3.1. *Let $\Omega \subset \mathbb{R}^{n+1}$ be a bounded, convex, open set with smooth boundary. Given $u_0 \in \mathbb{R}$ and $\alpha > 0$, suppose that $u \in C^1(\bar{\Omega})$ has a single critical point, $p \in \Omega$, is smooth in $\Omega \setminus \{p\}$, and satisfies*

$$\begin{cases} |Du|f^\alpha \left(-\frac{Du}{|Du|}, -\frac{1}{|Du|} \left[I - \frac{Du \otimes Du}{|Du|^2} \right] \cdot D^2 u \right) = 1 & \text{in } \Omega \setminus \{p\} \\ u = u_0 & \text{on } \partial\Omega, \end{cases}$$

where $f : S^n \times \Gamma_+^{n \times n} \rightarrow \mathbb{R}$ is monotone non-decreasing and inverse-concave. Then $w := ((1 + \alpha)(u - u_0))^{\frac{1}{1+\alpha}}$ is concave.

If we no longer assume that Ω is bounded, but require instead that $u_0 = -\infty$ and that the level sets of u are bounded, then u is concave.

Proof. Consider the concavity function $Z : [0, 1] \times \Omega \times \Omega \rightarrow \mathbb{R}$, which we recall is defined by

$$Z(r, x, y) := w(rx + (1 - r)y) - (rw(x) + (1 - r)w(y)).$$

Choose the triple (r_0, x_0, y_0) so that

$$Z(r_0, x_0, y_0) = \min_{[0,1] \times \bar{\Omega} \times \bar{\Omega}} Z(r, x, y).$$

As before, it suffices to assume that r_0 , x_0 and y_0 are interior points. Let us abuse notation by writing $Z(x, y) := Z(r_0, x, y)$. Then (x_0, y_0) is a stationary point of Z and hence, setting $z_0 := r_0x_0 + (1 - r_0)y_0$,

$$0 = \partial_{x^i} Z(x_0, y_0) = r_0(w_i(z_0) - w_i(x_0))$$

and

$$0 = \partial_{y^i} Z(x_0, y_0) = (1 - r_0)(w_i(z_0) - w_i(y_0)).$$

So

$$Dw(z_0) = Dw(x_0) = Dw(y_0) =: p_0.$$

We may also assume that $p_0 \neq 0$ since if $p_0 = 0$, we would have $x_0 = y_0 = z_0$, and hence $Z(x_0, y_0) = 0$.

At this point, the proof differs from that of previously known results. In order to obtain the best possible result, we need to optimize the second variation inequality for Z (cf. [5, 6, 7]). Since (x_0, y_0) is a local minimum, we obtain, for any pair of endomorphisms a and b of \mathbb{R}^{n+1} ,

$$\begin{aligned} 0 &\leq \left. \frac{d^2}{ds^2} \right|_{s=0} Z(x_0 + sa \cdot e_i, y_0 + sb \cdot e_j) \\ &= (a_i^p \partial_{x^p} + b_i^p \partial_{y^p})(a_j^q \partial_{x^q} + b_j^q \partial_{y^q}) Z(x_0, y_0) \\ &= (r_0 a + (1 - r_0) b)_i^p (r_0 a + (1 - r_0) b)_j^q w_{pq}(z_0) \\ &\quad - r_0 a_i^p a_j^q w_{pq}(x_0) - (1 - r_0) b_i^p b_j^q w_{pq}(y_0). \end{aligned}$$

The endomorphisms a and b will be chosen in order to optimize this inequality. Denote by

$$\pi_0 := I - \frac{p_0 \otimes p_0}{|p_0|^2}$$

the projection onto the orthogonal complement of p_0 . Since the equation is degenerate in the direction of Du , we consider only those endomorphisms of the form

$$a = \hat{a} \circ \pi_0 \quad \text{and} \quad b = \hat{b} \circ \pi_0,$$

where \hat{a} and \hat{b} are endomorphisms of $\pi_0 \cdot \mathbb{R}^{n+1}$. Then

$$(19) \quad \hat{c}_i^p \hat{c}_j^q (A_{z_0})_{pq} \leq r_0 \hat{a}_i^p \hat{a}_j^q (A_{x_0})_{pq} + (1 - r_0) \hat{b}_i^p \hat{b}_j^q (A_{y_0})_{pq},$$

where $\hat{c} := r_0 \hat{a} + (1 - r_0) \hat{b}$ and

$$A_x := -\frac{1}{|Dw(x)|} \left(I - \frac{Dw(x) \otimes Dw(x)}{|Dw(x)|^2} \right) \cdot D^2 w(x).$$

Setting $\hat{a} = A_{x_0}^{-1}$ and $\hat{b} = A_{y_0}^{-1}$, we find

$$r_0 A_{x_0}^{-1} + (1 - r_0) A_{y_0}^{-1} \leq A_{z_0}^{-1}.$$

The monotonicity and concavity of f_* then yield

$$\begin{aligned}
 w(z_0) &= |p_0|^{-\frac{1}{\alpha}} f^{-1} \left(\frac{p_0}{|p_0|}, A_{z_0} \right) \\
 &= |p_0|^{-\frac{1}{\alpha}} f_* \left(\frac{p_0}{|p_0|}, A_{z_0}^{-1} \right) \\
 &\geq |p_0|^{-\frac{1}{\alpha}} f_* \left(\frac{p_0}{|p_0|}, r_0 A_{x_0}^{-1} + (1 - r_0) A_{y_0}^{-1} \right) \\
 &\geq r_0 |p_0|^{-\frac{1}{\alpha}} f_* \left(\frac{p_0}{|p_0|}, A_{x_0}^{-1} \right) + (1 - r_0) |p_0|^{-\frac{1}{\alpha}} f_* \left(\frac{p_0}{|p_0|}, A_{y_0}^{-1} \right) \\
 &= r_0 w(x_0) + (1 - r_0) w(y_0).
 \end{aligned}$$

The first claim is proved. The second follows as in Theorem 2.1. \square

As a corollary, we obtain differential Harnack inequalities for flows by positive powers of inverse-concave speeds which contract convex hypersurfaces to points. Such inequalities were already observed by Andrews [2, Corollary 5.11] and Chow [9].

Corollary 3.2. *Let $\{\mathcal{M}_t^n\}_{t \in [t_0, T]}$ be a smooth family of boundaries $\mathcal{M}_t^n = \partial\Omega_t$ of bounded convex bodies Ω_t moving with inward normal speed*

$$F(x, t) = f^\alpha(\nu(x, t), [A_{(x,t)}])$$

for some $\alpha > 0$, where $f : S^n \times \Gamma_+^{n \times n} \rightarrow \mathbb{R}_+$ is a smooth function which is $\text{SO}(n)$ -invariant, monotone non-decreasing and inverse-concave in its second entry. Suppose that the hypersurfaces \mathcal{M}_t^n contract to a point at time T . Then the $(1 + \alpha)$ -th root $w := ((1 + \alpha)(u - t_0))^{\frac{1}{1+\alpha}}$ of the arrival time $u : \Omega_{t_0} \rightarrow \mathbb{R}$ is concave. Equivalently,

$$\partial_t F + 2\nabla_V F + A(V, V) + \frac{\alpha F}{(1 + \alpha)(t - t_0)} \geq 0 \text{ for all } V \in T\mathcal{M}_t, t \in (t_0, T).$$

If the solution is ancient, then u is concave. Equivalently,

$$\partial_t F + 2\nabla_V F + A(V, V) \geq 0 \text{ for all } V \in T\mathcal{M}_t, t \in (t_0, T)$$

Proof. The proof is similar to that of Corollary 2.2. \square

Remark 3.3. *Corollary 3.2 assumes that the solution contracts to a single point at the singular time. This is known to be the case for solutions to isotropic flows satisfying only slightly stronger conditions than α -inverse-concavity [8, Theorem 5]. (The proof of this fact does not require differential Harnack inequalities). Moreover, examples are given in [8] of speeds which do not preserve convexity of the level sets \mathcal{M}_t^n , and hence cannot admit power concave arrival times.*

We do not require that the limiting shape is round. Indeed, in many situations where Harnack inequalities are known, this will not be the case [3, 4].

In contrast to the known approaches to differential Harnack inequalities, Theorem 3.1 does not require any regularity hypotheses for the speed.

REFERENCES

- [1] ALVAREZ, O., LASRY, J.-M., AND LIONS, P.-L. Convex viscosity solutions and state constraints. *J. Math. Pures Appl. (9)* 76, 3 (1997), 265–288.
- [2] ANDREWS, B. Harnack inequalities for evolving hypersurfaces. *Math. Z.* 217, 2 (1994), 179–197.
- [3] ANDREWS, B. Contraction of convex hypersurfaces by their affine normal. *J. Differential Geom.* 43, 2 (1996), 207–230.
- [4] ANDREWS, B. Classification of limiting shapes for isotropic curve flows. *J. Amer. Math. Soc.* 16, 2 (2003), 443–459.
- [5] ANDREWS, B. Pinching estimates and motion of hypersurfaces by curvature functions. *J. Reine Angew. Math.* 608 (2007), 17–33.
- [6] ANDREWS, B., AND CLUTTERBUCK, J. Proof of the fundamental gap conjecture. *J. Amer. Math. Soc.* 24, 3 (2011), 899–916.
- [7] ANDREWS, B., AND CLUTTERBUCK, J. Sharp modulus of continuity for parabolic equations on manifolds and lower bounds for the first eigenvalue. *Anal. PDE* 6, 5 (2013), 1013–1024.
- [8] ANDREWS, B., MCCOY, J., AND ZHENG, Y. Contracting convex hypersurfaces by curvature. *Calc. Var. Partial Differential Equations* 47, 3-4 (2013), 611–665.
- [9] CHOW, B. On Harnack’s inequality and entropy for the Gaussian curvature flow. *Comm. Pure Appl. Math.* 44, 4 (1991), 469–483.
- [10] CHOW, B., AND CHU, S.-C. Space-time formulation of Harnack inequalities for curvature flows of hypersurfaces. *J. Geom. Anal.* 11, 2 (2001), 219–231.
- [11] COLDING, T. H., AND MINICOZZI, II, W. P. Differentiability of the arrival time. *Comm. Pure Appl. Math.* 69, 12 (2016), 2349–2363.
- [12] EVANS, L. C., AND SPRUCK, J. Motion of level sets by mean curvature. I. *J. Differential Geom.* 33, 3 (1991), 635–681.
- [13] HAMILTON, R. S. Harnack estimate for the mean curvature flow. *J. Differential Geom.* 41, 1 (1995), 215–226.
- [14] HELMENS DORFER, S., AND TOPPING, P. The geometry of differential Harnack estimates. arxiv:1301.1543, 2013.
- [15] HUISKEN, G. Flow by mean curvature of convex surfaces into spheres. *J. Differential Geom.* 20, 1 (1984), 237–266.
- [16] HUISKEN, G. Local and global behaviour of hypersurfaces moving by mean curvature. In *Differential geometry: partial differential equations on manifolds (Los Angeles, CA, 1990)*, vol. 54 of *Proc. Sympos. Pure Math.* Amer. Math. Soc., Providence, RI, 1993, pp. 175–191.
- [17] JUUTINEN, P. Concavity maximum principle for viscosity solutions of singular equations. *NoDEA Nonlinear Differential Equations Appl.* 17, 5 (2010), 601–618.

- [18] KAWOHL, B. *Rearrangements and convexity of level sets in PDE*, vol. 1150 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1985.
- [19] KENNINGTON, A. U. Power concavity and boundary value problems. *Indiana Univ. Math. J.* *34*, 3 (1985), 687–704.
- [20] KOREVAAR, N. J. Convex solutions to nonlinear elliptic and parabolic boundary value problems. *Indiana Univ. Math. J.* *32* (1983), 603–614.
- [21] KOTSCHWAR, B. Harnack inequalities for evolving convex surfaces from the space-time perspective. math.la.asu.edu/~kotschwar/pub, 2009.
- [22] SAKAGUCHI, S. Concavity properties of solutions to some degenerate quasilinear elliptic Dirichlet problems. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)* *14*, 3 (1987), 403–421 (1988).
- [23] TRUDINGER, N. S. The Dirichlet problem for the prescribed curvature equations. *Arch. Rational Mech. Anal.* *111*, 2 (1990), 153–179.
- [24] WANG, X.-J. Convex solutions to the mean curvature flow. *Ann. of Math. (2)* *173*, 3 (2011), 1185–1239.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TENNESSEE KNOXVILLE,
KNOXVILLE TN, 37996-1320

E-mail address: `tbourni@utk.edu`

E-mail address: `mlangford@utk.edu`

SCHOOL OF MATHEMATICAL AND PHYSICAL SCIENCES, THE UNIVERSITY OF
NEWCASTLE, NEWCASTLE, NSW, AUSTRALIA, 2308

E-mail address: `mathew.langford@newcastle.edu.au`